

Why Are Radio-Galaxies Prolific Producers of Type Ia Supernovae?[†]

Massimo Della Valle

Istituto Nazionale di Astrofisica, Osservatorio Astrofisico di Arcetri,

Largo Enrico Fermi 5, I-50125 Florence, Italy

Nino Panagia¹

Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218

Paolo Padovani

European Southern Observatory, Karl-Schwarzschild-Strasse 2, Garching bei München,

Germany

Enrico Cappellaro

Istituto Nazionale di Astrofisica, Osservatorio Astronomico di Capodimonte

Salita Moiariello, 16, I-80131, Napoli, Italy

Filippo Mannucci

CNR-IRA

Largo Enrico Fermi 5, I-50125 Florence, Italy

Massimo Turatto

Istituto Nazionale di Astrofisica, Osservatorio Astronomico di Padova,

Vicolo Osservatorio 5, Padova, Italy

Received _____; accepted _____

submitted to the Ap.J.

[†]**Version 6 of February 2, 2008**

¹On assignment from the Space Telescope Operations Division, Research and Scientific Support Department of ESA

ABSTRACT

An analysis of SNIa events in early type galaxies from the Cappellaro *et al.* (1999) database provides conclusive evidence that the rate of type Ia Supernovae (SNe) in radio-loud galaxies is about 4 times higher than the rate measured in radio-quiet galaxies, *i.e.* $\text{SNIa-rate}(\text{radio} - \text{loud galaxies}) = 0.43_{-0.14}^{+0.19} h_{75}^2$ SNu as compared to $\text{SNIa-rate}(\text{radio} - \text{quiet galaxies}) = 0.11_{-0.03}^{+0.06} h_{75}^2$ SNu. The actual value of the enhancement is likely to be in the range $\sim 2 - 7$ ($P \sim 10^{-4}$). This finding puts on robust empirical grounds the results obtained by Della Valle & Panagia (2003) on the basis of a smaller sample of SNe.

We analyse the possible causes of this result and conclude that the enhancement of SNIa explosion rate in radio-loud galaxies has the same origin as their being strong radio sources, but there is no causality link between the two phenomena. We argue that repeated episodes of interaction and/or mergers of early type galaxies with dwarf companions, on times-scale of about 1 Gyr, are responsible for inducing both strong radio activity observed in $\sim 14\%$ of early type galaxies and to supply an adequate number of SNIa progenitors to the stellar population of ellipticals.

Subject headings: (stars:) stars: supernovae – galaxies: radio

1. Introduction

The Supernova rate is a key quantity for astrophysics. Its knowledge is essential to constrain the mechanisms of galaxy formation and for understanding galaxy evolution. In particular, the chemical evolution of systems that at the present epoch have substantially reduced or totally exhausted star formation activity, such as elliptical galaxies, may dramatically depend on the rate of type Ia events (e.g. Renzini et al. 1993, Maoz & Gal-Yam 2004). Thus, any mechanism capable of producing a substantial enhancement of SNIa rate in this type of galaxies should be carefully investigated. Moreover, studies of the connection between the frequency of occurrence of type Ia SNe and the properties of the environment where the SNe-Ia originate (e.g. Della Valle & Livio 1994; Mannucci et al. 2005) can effectively constraint the still controversial nature of their progenitors (e.g. Hamuy et al. 2003, Livio and Riess 2003).

In recent years three works have dealt with this topic, namely Capetti (2002), Livio, Riess and Sparks (2002) and Della Valle and Panagia (2003). The former two papers claim the existence of a *causality* link between the position of 6 type Ia SNe (Capetti) and a dozen of nova stars (Livio et al.) with the jets of the respective parent radio-galaxies. In this scenario it is the jet itself that modifies the physical conditions of its environment by increasing the accretion rate from the interstellar medium onto the white dwarf progenitors, and thus it is the *direct cause* of a local enhancement of Nova and type Ia SN explosions.

Della Valle & Panagia, after analysing the Evans, van den Bergh & Clure (1989) database (hereafter EVC89), provided strong evidence for the existence of an enhancement by a factor 4–15 of the SN-Ia rate in radio-loud galaxies with respect to radio-quiet ones. They argued that the mechanism which boosts the SN production is essentially independent of the position of the Supernovae inside the parent galaxy, rather having the same common origin as the parent galaxies being strong radio sources, *without causality* link between the

two phenomena. They suggested that repeated episodes of interaction and/or mergers of early type galaxies with dwarf companions are responsible for inducing both strong radio activity in $\sim 14\%$ of early type galaxies and the ~ 1 Gyr old stellar population needed to supply an adequate number SNIa progenitors. However, their results were based on a relatively small sample and deserve further investigation.

In this paper we analyse an approximately five times larger sample of SNe occurred in early type galaxies, as reported by Cappellaro, Evans & Turatto (1999). This sample was obtained by merging the Cappellaro et al. (1993, hereafter C93) sample with EVC89. In § 2 we present the dataset, in § 3 we first analyse the C93 sample, with the aim of providing an independent check of the results obtained by Della Valle & Panagia (2003) (based on EVC89), then we compute the SN rate in radio-loud and radio-quiet galaxies on the whole sample of SNe provided by Cappellaro et al. (1999). In § 4 we discuss the results and their implications on the nature of SNIa progenitors.

Tab. 1 gives the comparison between the EVC89 and Cappellaro et al. (1999) samples. Data for the C93 sample can be obtained by subtracting the EVC89 data from Cappellaro et al. (1999).

In the following, we will denote as *radio-loud* sources, all galaxies which have radio luminosities at 1.4GHz higher than $10^{29} \text{ erg s}^{-1} \text{ Hz}^{-1}$ (e.g. Sadler, Jenkins, & Kotanyi 1989, Ledlow & Owen 1996). The threshold value of $\sim 10^{29} \text{ erg s}^{-1} \text{ Hz}^{-1}$ is the faint limit of the luminosity function for the radio-galaxies (see also Urry & Padovani 1995), and, therefore, represents the ‘natural’ threshold to identify “radio-loud” galaxies. We also define *radio-faint* the galaxies characterized by $4 \times 10^{27} \leq P \leq 10^{29} \text{ erg s}^{-1} \text{ Hz}^{-1}$, i.e. galaxies whose radio emission is still not dominated by thermal processes (Phillips et al. 1986; and Sadler et al. 1989) and finally *radio-quiet* all galaxies with radio luminosities below $P = 4 \times 10^{27} \text{ erg s}^{-1} \text{ Hz}^{-1}$.

2. Statistics of SNIa explosions in early type galaxies

The database compiled by Cappellaro et al. (1999) has the advantage of being defined ‘a priori’, i.e. the monitored galaxies were selected before the SN explosions were discovered. As a consequence our statistical analysis will not suffer of any kind of ‘a posteriori’ selection criteria, which actually may affect the statistical analysis carried out on catalogues of parent galaxies of SNe. We have first obtained from the NASA/IPAC Extragalactic Database (NED) the most accurate positions for the sources in the Cappellaro et al.’s database, restricting ourselves to morphological Hubble types $T \leq -1.5$. We then cross-correlated this list with the NRAO VLA Sky Survey (NVSS; Condon et al. 1998), which covers the whole sky north of -40° at 1.4 GHz. For galaxies below this limit we used the Parkes-MIT-NRAO survey (PMN; Griffith & Wright 1993) at 4.85 GHz, converting the flux to 1.4 GHz assuming a radio spectral index equal to -0.75 . Sources without an NVSS radio match were dealt with in two steps. First, to take care of cases where the NVSS might be resolving out the source or where the core was very weak and most of the flux was coming from the lobes (which could be way off the optical position), we cross-correlated our database with the PMN (in the south) and the GB6 (Gregory et al. 1996; in the north) catalogues. Second, if still no match was found, we assumed a radio flux < 2.5 mJy, the limit of the NVSS. The adopted correlation radius was chosen according to the uncertainties on the optical and radio positions of the sources. Namely, a radius of 20 and 30'' was taken for the NVSS and PMN/GB6 respectively for galaxies having optical positional uncertainties $< 5''$. Radii of 30 and 45'' were adopted for galaxies having larger positional errors. Note that this approach is conservative as it might still miss some interacting galaxies for which the optical position could be quite different from the radio one. The NVSS will also underestimate the total flux for extended radio sources. This cannot have a major effect on our results for two reasons: 1. we checked all of the galaxies

in Tab. 2 and found only one such case, NGC 1316, whose classification did not change when using the total flux; 2. we find a fraction of radio-loud sources $\sim 12\%$ (see below), perfectly consistent with the value generally quoted (Ledlow & Owen 1996). Finally we note that the offset of $\sim 20''$ between the optical and the PMN position of E471–G27 is consistent with the PMN positional uncertainty. However, the optical and NVSS positions differ by $\sim 1.2'$. Given that NED includes amongst this source’s names PMN J2351–2758 and TXS 2349–282 and that the NVSS image shows that the optical source lies at the edge of the radio contours, we decided to associate a radio source with this galaxy.

Out of a total of 2217 galaxies we have found 267 radio-loud, 212 radio-faint, and 1738 radio-quiet galaxies respectively. The twenty-one galaxies in which SNe were discovered are reported in Tab. 2. Col. 1 and 2 give the SN and the galaxy designation, col. 3 the morphological Hubble type, col. 4 the radio flux and col. 5 the rate of decline of the luminosity expressed as Δm_{15} (see Phillips 1993). Fig. 1 shows the frequency distribution of all 2215 early type galaxies belonging to our sample as a function of their absolute B-band magnitudes, M_B (solid line). The dashed line refers to the distribution of the general luminosity function of ellipticals (adapted from Muriel et. al. 1995); the shaded histogram denotes the distribution of radio-loud galaxies. The absolute magnitudes of the galaxies have been taken from the RC3 catalogue (de Vaucouleurs et al. 1991). In tab. 1 we report the galaxy type (col. 1), the number of galaxies for each type used by Della Valle & Panagia (2003) and in this paper (col. 2 and col. 5), the normalized control times, *i.e.* the product of the control time (*i.e.* the time during which a given galaxy has been observed) to the galaxy’s B-band absolute luminosity, so that is given in units of $\text{years} \times 10^{10} L_{\odot}^B$ (col. 3 and col. 6), and the number of type Ia SNe discovered (col. 4 and col. 7). $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$ is adopted throughout the paper.

An inspection to Fig.1 and Tab.1 reveals the following:

i) The Cappellaro et al. (1999) sample is not biased in favour of bright galaxies. It shows indeed a deficiency in the number of galaxies brighter than $M_B = -21$ in comparison with a typical luminosity function of early type galaxies (e.g. Muriel, Nicotra & Lambas 1995). As apparent from Fig. 1 the distribution of the monitored galaxies has not the marked deficiency of galaxies fainter than $M_B = -19$ which characterized the ‘naked-eye’ Evans sample.

ii) The fraction of radio-loud galaxies in our sample is 0.12 ± 0.02 of the whole sample of galaxies. Such fraction is essentially identical to the value 0.140 ± 0.024 as reported by Ledlow & Owen (1996) for radio galaxies having the same radio luminosities ($> 10^{29}$ erg s^{-1} Hz $^{-1}$ at 1.4GHz). This indicates that the frequency of radio galaxies in our sample is not affected by biases related to their radio properties.

iii) Considering all early type galaxies, regardless of their radio properties, an overall SNIa rate of $0.19^{+0.05}_{-0.04} h_{75}^2$ SNU 2 is obtained.

3. Analysis and Results

Since the Cappellaro et al. (1999) sample includes the EVC89, we have first carried out an independent analysis of the results obtained by Della Valle & Panagia, just using C93 (which does not include EVC89). Separating the galaxies into radio-loud and radio-quiet (this subclass includes also radio-faint for homogeneity with the previous paper) we obtain 5.5 and 11.5 SNIa events, respectively. Non-integer numbers derive from NGC 3106 which is a border line object between the two classes having a radio luminosity of $\sim 9.2 \times 10^{28}$

$^2 1\text{SNU} = 1 \text{ SN}(100\text{yr})^{-1} (10^{10} L_{\odot}^B)^{-1}$

erg/s/Hz, and therefore we have attributed 0.5 SN to each bin³ The SN rate is derived as a ratio between the number of detected SNe and the respective controltime (expressed in units of 100 years) and it gives: $r_{loud} = 0.41^{+0.25}_{-0.20}h_{75}^2\text{SNu}$ and $r_{quiet} = 0.17^{+0.07}_{-0.05}h_{75}^2\text{SNu}$. The errors corresponds to 1σ and have been computed from a direct application of Poisson statistics in the small number regime (e.g. Ebeling 2003, Gehrels 1986). A *T-Student* test rejects the null hypothesis between radio-loud and radio-quiet galaxy samples with a probability of $P \lesssim 5 \times 10^{-3}$. This result points out the high statistical significance of C93 sample itself and provides independent evidence for the existence of an overproduction of SNe-Ia in radio-loud galaxies. With this in mind we can now merge the EVC89 and C93 samples and determine the rate of type Ia SNe in radio-galaxies on even more robust empirical grounds. The enlarged sample has allowed us to allocate 3 classes of galaxies on the basis of their radio activity, as pointed out in Section 1. We have 9.5, 4 and 7.5 SNIa events in radio-loud, radio-faint and radio-quiet galaxies, respectively. Fractional numbers derive from two border line objects, NGC 3106 (see above) and NGC 4526 which radio luminosity is $\sim 4 \times 10^{27}$ erg/s/Hz. For the SN rates we obtain: $r_{loud} = 0.43^{+0.19}_{-0.14}h_{75}^2\text{SNu}$, $r_{faint} = 0.23^{+0.18}_{-0.11}h_{75}^2\text{SNu}$ and $r_{quiet} = 0.11^{+0.06}_{-0.03}h_{75}^2\text{SNu}$, respectively. The errors corresponds to 1σ and have been computed as above. A *T-Student* test rejects the null hypothesis between radio-loud and radio-quiet galaxy samples with a probability of $P \sim 4 \times 10^{-4}$.

The comparison among the SN rates, normalized to the B luminosity, in different Hubble types galaxies should be taken with some degree of caution because the B luminosity

³Attributing half SN event to each bin for objects close to the luminosity threshold may appear a quite unusual procedure, nevertheless it has been adopted to minimize the effects due to the uncertainty on H_0 on the computation of the radio power. This approach may affect the measurements of the SN rates in radio -loud and -quiet galaxies up to $\sim 6\%$ and 10% , respectively.

is a poor tracer of the galaxy mass in stars. In this case, since our analysis is restricted to E/S0 Hubble types ($T \leq -1.5$), this approximation is still acceptable (see Tab. 3). However, the recent release of the Two Micron All Sky Survey (2MASS) collaboration (Jarret et al. 2003), allows us to normalize the rates to the IR luminosity (K-band) and subsequently to the mass in stars of the parent galaxies by fitting spectrophotometric models (e.g. Brinchmann & Ellis 2000; Dickinson et al. 2003) through the broad band fluxes (see Mannucci et al. 2005 for details). In Tab. 4 we also report the SN rate per unit of stellar mass.

4. Discussion and Conclusions

Our results allow us to confidently conclude that the SNIa rate in radio-loud galaxies is a full factor of $4.4^{+2.4}_{-2.6}$ higher than it is measured in radio-quiet objects confirming, on more robust statistical grounds, the findings of Della Valle & Panagia (2003). We have also obtained some evidence ($P \lesssim 5 \times 10^{-2}$) that the class of radio-faint galaxies may have a rate that is intermediate between radio-loud and radio-quiet objects. This enhancement in SNIa events in radio galaxies can be explained in basically two different ways:

1) The enhanced SNIa explosion rate in radio-loud galaxies is a *direct* consequence of their being strong radio sources. This is the scenario suggested by Capetti (2002) and Livio et al. (2002). According to these authors the accretion rate from the interstellar medium onto white dwarfs could be enhanced by the action of the jets up to either driving an accreting white dwarf to approach the Chandrasekhar mass and trigger a SNIa explosion (Livio 2000, and references therein) or increasing the accretion rate from the interstellar medium onto the white dwarfs by 1-2 orders of magnitude, up to $\sim 10^{-8/-9} M_{\odot} \text{ yr}^{-1}$, which is the rate necessary to trigger classical nova explosions (Priyalnik & Kovetz 1995). This

process might account for the high rate of novae observed in M87 in the regions nearby the jets. However, we note that equation (4) of Livio et al (2002), which is the transcription of Bondi’s (1952) formula for accretion on a star at rest, is valid only for very low stellar velocities ($\ll 1 \text{ km s}^{-1}$). This fact has important consequences on the practicability of this scenario. In a rather optimistic scenario in which $v_{star} = 1 \text{ km s}^{-1}$ and the jet has a width of $\sim 100 \text{ pc}$, the transit time is about 100 Myrs during which the total accreted mass would be less than $0.1 M_{\odot}$, i.e. hardly enough to make a WD explode as SNIa, unless its mass was greater than $\sim 1.35 M_{\odot}$. In a more realistic case, in which one assumes typical star velocities of the order of $10\text{-}100 \text{ km s}^{-1}$ (see for example Williams 1994, who has measured the velocities of nova systems in the Milky Way) the accreted mass onto massive WDs would be only $10^{-5}\text{-}10^{-8} M_{\odot}$. This makes the accretion process from ISM highly inefficient to trigger SNIa explosions (while it may play a not negligible role to trigger nova explosions) even in the presence of a high compression regime. In addition, we note that:

i) Even under the best conditions that could induce a SNIa explosion ($v \ll 1 \text{ km s}^{-1}$; $M_{WD} \sim 1.2 \div 1.3 M_{\odot}$), the resulting SNIa is expected to be spatially confined to the regions immediately adjacent to radio jets and/or the bulk of radio activity. While this may be true for NGC 4374 (Fig. 2), it appears not to be the case for Fornax A=NGC 1316 (Fig. 3) in which the two SNIa (SN 1980N and SN 1981D) are located quite far from the strong radio lobes (Geldzahler & Fomalont 1984), nor for Centaurus A=NGC 5128, in which SN 1986G is deeply embedded in the equatorial lane of gas and dust (Fig. 4).

ii) If the dominant phenomenon is accretion from the ISM rather than from a binary companion, an extreme consequence is that all *single* WDs could contribute to the Nova and SNIa production, and not only some binary systems with suitable parameters. Actually, since the fraction of stars that may give rise to a SNIa is only $\eta \sim 5 - 10\%$ of the total number of stars with masses above $3 M_{\odot}$ (Madau, Della Valle & Panagia 1998, Dahlen et al. 2004), in order to explain the observed enhancement in the SN rate (a factor ~ 4) one

can work out that jets should be capable of raising the efficiency of the SN explosions up to $\sim 100\%$ not only in the regions near the jets, but also over a significant fraction of the entire galaxy, according to the simple relation: $\eta_{jet}/\eta_{gal} \sim 3 \times vol_{gal}/vol_{jet}$ where η parameter is the efficiency of SN explosions and vol_{jet} and vol_{gal} are the volumes sampled by the jet and the whole galaxy.

iii) For a density of 100 cm^{-3} and a jet width of 100 pc as advocated by Livio et al. (2002), the column density of HI associated with a jet would be about $3 \times 10^{22} \text{ cm}^{-2}$. In this case one should expect an intense 21cm line emission, which has never been observed in association with jets.

iv) A clear indication that accretion of interstellar matter by slow ($v < 1 \text{ km s}^{-1}$) WDs is not an efficient process to trigger SNIa explosions, is that SNIa occurring in spiral galaxies are not associated with molecular clouds, which are regions with the “ideal” conditions, i.e. high particle density and very low temperatures, for triggering a SNIa explosion through this mechanism. It appears that the accretion from ISM can work only in the presence of very special circumstances, which makes a SNIa explosion through accretion “à la Bondi” an exception rather than the rule.

In summary, since there is no evidence that direct accretion from the ISM could be capable to boost the SNIa rate in radiogalaxies by a factor of 4, we conclude that the dominant process is very likely a different one.

2) As an alternative we propose that the enhancement of SNIa explosion rate in radio-loud galaxies has the *same common* origin as their being strong radio sources (see also Kochhar 1989; Kochhar & Prabhu 1984), but that there is no *causality* link between the two phenomena. If the radio activity of a galaxy is triggered by interaction and/or mergers (e.g. Baade & Minkowski 1954, Balick & Heckman 1982, Heckman *et al.* 1986), then the SNIa rate can be enhanced by the formation or the capture of relatively young stellar populations in which SNIa occur at much higher rates than in genuinely old populations

(Della Valle & Livio 1994, Panagia 2000, Mannucci et al. 2005). SNIa progenitors are believed to be stars with original masses $\gtrsim 3 M_{\odot}$ and $\lesssim 8 M_{\odot}$ (Greggio & Renzini 1983) having nuclear-burning lifetimes shorter than ~ 400 Myrs. After that time, the star becomes a white dwarf and may explode as a SNIa only after an additional time as needed for either accreting mass from a companion or merging with it. Such a time (τ) is essentially unconstrained by theory, but measurements of SNIa rates in the local universe (Mannucci et al. 2005) have shown that: (a) the rate of type Ia SNe in blue galaxies is about a factor 30 higher than it has been measured in red galaxies; (b) the rate of type Ia SNe can be reproduced by combining a constant contribution (corresponding to the rate of Ia in radio-quiet galaxies) + another contribution which exhibits a behaviour similar to that of core-collapse SNe (see their Fig. 6). This implies evolutionary times (main sequence life-time + $\tau \lesssim 0.2$ Gyrs) much shorter than the time scale needed by young ‘blue’ galaxies to become ‘red’ galaxies (~ 0.8 Gyrs to change (B-K) by about 1 mag, see for example Bruzual & Charlot 2003). This result may not be at variance with the recent findings of Gal-Yam & Maoz (2004) and Stogler et al. (2004), who derived $\tau \sim 2 - 4$ Gyrs from counts of SNeI-a at high z . This may rather suggest that τ spans a broad range of values, possibly from $\lesssim 0.2$ up to a few Gyrs (Mannucci, Della Valle & Panagia, in preparation, Greggio 2005). As a consequence, in order to sustain a relatively high rate of SNIa explosions (characterized by a short τ) during the time in which a galaxy is radio-loud ($\Delta T_{RL} \sim 10^8$ yr, e.g. Srianand & Gopal-Krishna 1998), a galaxy has to have a new injection of young stellar populations at suitable time intervals. Since the fraction of radio-loud ellipticals is about 14% of the total number of ellipticals (Ledlow & Owen 1996), we can estimate that, on average, an elliptical galaxy undergoes a number of star formation episodes, n_{inj} , of the order of $n_{inj} \simeq 0.14 \times T_H / \Delta_{RL} \sim 20$, for a Hubble time $T_H \simeq 14$ Gyrs. Therefore, the time interval between successive episodes of injection of young stellar populations, Δt_{inj} , turns out to be $\Delta t_{inj} = T_H / n_{inj} \simeq 0.7$ Gyr. In late type galaxies, the ongoing

star formation provides the needed input for a steady production of SNIa events. Such a supply in early type galaxies can naturally be provided by repeated episodes of interaction or mergers that induces either formation of young stellar populations (galaxy interaction; *e.g.* the Antennae galaxies, Whitmore *et al.* 1999) or capture of young stars from dwarf companions. Well known examples of interacting and/or merging early type galaxies are Centaurus A=NGC 5128 and Fornax A=NGC 1316, which are strong radio sources and indeed have produced 1+2 SNIa in the last century.

Strong support to this hypothesis comes from the following two facts:

i) In Tab. 4 we have reported the SN rate for Ellipticals radio- loud, faint, quiet, and Spirals normalized per unit of mass. A simple glance at col. 2 and Fig. 5 shows that the rates of SNIa are higher in late type galaxies than in early type galaxies by a full order of magnitude, then suggesting a connection between the production of type Ia SNe and star formation activity (*e.g.* Mannucci *et al.* 2005). Fig. 5 shows the behaviour of the normalized SNIa rate (per $10^{10} M_{\odot}$) as a function of the morphological Hubble type and radio activity. The data clearly shows that for the SNIa production, radio-loud early type galaxies behave like ‘S0a/b’ which are systems where the star forming activity is certainly dimmed with respect to Sd/Irr, but still active, likely of the order of $0.1 M_{\odot} \text{ yr}^{-1}$ per $10^{10} M_{\odot}$ (*e.g.* Tinsley 1974, Sandage 1986, Kennicutt 1998). It is then plausible that this enhancement of SNIa rate originates from episodes of star formation, which may have been triggered, on time-scale of 1 Gyr, by merging dwarf galaxies. This scenario provides a natural explanation of why among early type galaxies SNIa rates are higher in radiogalaxies without requiring the concurrence of any new and/or additional process to enhance the SNIa rates.

ii) The existence of systematic differences between type Ia SNe occurring in Spirals and Ellipticals, which has been accumulating in the last decade both spectroscopically

(Branch & van den Bergh 1993, Li et al. 2001) and photometrically (van den Bergh & Pazder 1992, Della Valle & Panagia 1992, Hamuy et al. 1995, Hamuy et al. 2000), can be effectively represented by the distribution of the decline rate parameter Δm_{15} (Phillips 1993; Phillips et al. 1999). In Fig. 6 we report the distribution of the rates of decline for late (shade-dotted histogram) and early type galaxies. Data were taken from Branch, Romanishin and Baron (1996) and Parodi et al. (2000). The probability that the two distributions originate from the same population, is excluded by KS-test at the level of $P=5 \times 10^{-7}$ (see also Altavilla et al. 2004). The most straightforward interpretation of the double-peaked distribution of the rates of decline is obtained in terms of an age sequence, (e.g. Branch & van den Bergh 1993; Ruiz-Lapuente 1995; Hamuy et al. 1996; Howell 2001; but see the case of SN 2002bo, Benetti et al. 2003), with metallicity being excluded or at least less important (Ivanov, Hamuy and Pinto 2000). With the dashed-solid box we have indicated the rates of decline for the 8 objects belonging to our sample of SNe in radio-galaxies for which it was possible to derive the rates of decline. We find tantalizing evidence ($P \lesssim 5 \times 10^{-2}$) that the rates of decline for the objects occurred in radio-galaxies are ‘intermediate’ between those of the late and early type galaxies. This fact supports the existence of a *continuum* of properties between galaxies with very active (Irr and late Spirals), active (early Spirals and radio-active Ellipticals) and exhausted (Ellipticals radio-quiet) star formation rate and type Ia supernovae. In Spirals with high star forming activity SNIa are generally overluminous, believed to be produced in younger systems from the most massive SNIa progenitors (*e.g.* Tutukov & Yungelson 1994, 1996, Ruiz-Lapuente, Burkert & Canal 1995), SNeIa in radio-loud ellipticals are characterized by faster rates of decline (*i.e.* less luminous SNe) and they might originate from progenitor of intermediate mass. Finally radio-quiet ellipticals mainly produce faint type Ia events from a genuine ‘old’ Pop II stellar population (light progenitors). An important consequence of our scenario is the predicted occurrence of a few type II-Ib/c SNe in early type galaxies at the early stages

of the same star formation burst that provides a steady supply of SNIa progenitors. Since core collapse SNe are produced by progenitors more massive than $8M_{\odot}$, whose lifetime is shorter than about 30 Myrs, only for a small fraction of the time between subsequent interaction/capture episodes (approximately $30\text{Myrs}/1\text{Gyr}=3\%$) would one expect to see such SNe in early type galaxies. Since in galaxies with active star formation the rate of type II and Ib/c SNe is about 2 and 2/3 times than that of SNIa, respectively, we estimate that one should detect core-collapse (*CC*) events at a rate $3\%\times 8/3 \simeq 8\%$ of SNIa events, which corresponds to an average rate of *CC* SNe of about $0.0152h_{75}^2\text{SNU}$. Given the control time measured by Cappellaro et al. (1999) for type *II plateau*, *II linear* and *Ib/c*: 4021 yr, 3213 yr and 5011 yr $\times L^B/10^{10}L_{\odot}^B$, and the relative frequencies of occurrence of each *CC* SN into the *plateau*, *linear* and *Ib/c* sub-types (0.6, 0.2 and 0.2 respectively), in our sample one should have detected $0.0152 \times (4057)/100=0.62$ SNe, where we have used the weighted mean of the control times reported above. Applying Poisson statistics, we estimate that the probability of finding no *CC* SN event in our sample is 54%, so that the absence of *CC* SNe is just a rather natural occurrence. We conclude that our prediction will be observationally testable when the SNIa statistics will have increased the control time by a factor ~ 6 ($\sim 2\sigma$ confidence level).

Our main conclusion is that the type Ia Supernova rate for radio-loud galaxies is more than a factor of four higher than for radio-quiet ones. This is true even when the rate is normalized to stellar mass instead of B-band luminosity, a more physical approach. Combined with the well-established connection between type Ia SNe and star-formation rate, this suggests a link between the radio-loud phenomenon and increased star-formation, which might prove to be an important piece of the still unexplained puzzle of radio-loudness in the Universe.

5. Acknowledgments

MDV is grateful to the Space Telescope Science Institute for the friendly hospitality and creative atmosphere. This research has made use of data from the NASA/IPAC Extragalactic Database (NED), which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. The authors wish to thank the referee Alessandro Capetti for his valuable comments, which have helped to improve the presentation of this paper.

Table 1: SN samples

galaxies	Evans			Cappellaro		
	N	Control Time	SNe	N	Control Time	SNe
radio-quiet	178	2270	0	1738	7127	7.5
radio-faint ¹	–	–	–	212	1770	4.0
radio-loud	19	847	4	267	2199	9.5
total	197	3117	4	2217	11096	21

¹Due to the scanty statistic the *radio-faint* subclass was not adopted in our previous paper (Della Valle & Panagia 2003). Radio-faint galaxies were included in the radio-quiet subclass.

Table 2: SNe occurred in early type galaxies belonging to the sample

SN	Galaxy	T	radio power	Δm_{15}
	NGC		$10^{29} \text{erg/s Hz}^{-1}$	
1961H	4564	-4.9	< 0.008	
1968A	1275	-1.6	720	
1970J	7619	-5.0	0.54	1.30
1972J	7634	-2.0	< 0.04	
1980I	4374	-4.7	20	
1980N	1316	-1.9	1000	1.28
1981D	1316	-1.9	1000	1.33
1981G	4874	-4.0	24	
1983G	4753	-2.2	0.53	1.50
1983J	3106	-1.9	0.92	
1986G	5128	-2.2	18	1.73
1990M	5493	-2.1	< 0.04	
1991Q	4926A	-1.6	0.31	
1991bg	4374	-4.7	20	1.88
1991bi	5127	-4.9	8.7	
1992A	1380	-1.9	< 0.02	
1992bo	E352-G57	-1.5	< 0.16	
1993C	2954	-4.9	< 0.08	
1993ah	E471-G27	-2.0	41	1.30
1994D	4526	-2.0	0.042	1.27
1996X	5061	-5.0	< 0.03	

Table 3: SN rates in units of SNe per century per $10^{10} L_{\odot}^B$

	<i>rate</i>	1σ	2σ	3σ
radio-loud	0.43	$+0.19$ -0.14	$+0.38$ -0.23	$+0.60$ -0.30
radio-faint	0.23	$+0.18$ -0.11	$+0.36$ -0.16	$+0.59$ -0.19
radio-quiet	0.11	$+0.06$ -0.03	$+0.11$ -0.06	$+0.18$ -0.07

Table 4: SN rates in units of SNe per century per $10^{10} M_{\odot}$

	<i>rate</i>	1σ	2σ	3σ
radio-loud	0.100	$+0.044$ -0.032	$+0.089$ -0.054	$+0.141$ -0.070
radio-faint	0.052	$+0.041$ -0.025	$+0.082$ -0.038	$+0.135$ -0.046
radio-quiet	0.023	$+0.012$ -0.008	$+0.024$ -0.013	$+0.035$ -0.018
all E's	0.044	$+0.016$ -0.014		
S0a/b	0.063	$+0.027$ -0.025		
Sbc/d	0.170	$+0.068$ -0.063		
Irr	0.77	$+0.42$ -0.31		

REFERENCES

- Altavilla, G., Fiorentino, G., Marconi, M. et al. 2004, MNRAS, 349, 1344
- Baade, W., & Minkowski, R. 1954, ApJ, 119, 206
- Balick, B., Heckman, T.M., 1982, ARA&A, 20, 431
- Benetti, S., Meikle, P., Stehle, M. et al. 2004, MNRAS, 348, 261
- Bondi, H. 1952, MNRAS, 112, 195
- Branch, D., & van den Bergh, S. 1993, AJ, 105, 2231
- Branch, D., Romanishin, W., & Baron, E. 1996, 470, L7
- Bruzual, G. & Charlot, S. 2003, MNRAS, 344, 1000
- Brinchmann, J., & Ellis, R.S. 2000 ApJ, 536, L77
- Capetti, A., 2002, ApJ, 574, L25
- Cappellaro, E., Turatto, M., Benetti, S., Tsvetkov, D. Yu., Bartunov, O. S., & Makarova, I. N 1993, A&A, 268, 472
- Cappellaro, E., Evans, R., & Turatto, M. 1999, A&A, 351, 459
- Condon, J. J., Cotton, W. D., Greisen, E. W., Yin, Q. F., Perley, R. A., Taylor, G. B., & Broderick, J. J. 1998, AJ, 115, 1693
- Dahlen et al. 2004, ApJ, 613, 189
- de Vaucouleurs G., de Vaucouleurs A., Corwin H.G., et al., 1991, Third Reference Catalogue of Bright Galaxies. Springer-Verlag, New York [RC3]
- Della Valle, M., & Livio, M., 1994, ApJ, 423, L31

- Della Valle, M., & Panagia, N., 1992, AJ, 104, 696
- Della Valle, M. & Panagia, N. 2003, ApJ, 587, L71
- Dickinson, M., Papovich, C., Ferguson, H. C., & Budavari, T. 2003, ApJ, 587, 25
- Ebeling, H. 2003, MNRAS, 340, 1269
- Evans, R, van den Bergh, S., & Clure, R.D., 1989, ApJ, 345, 752
- Gal-Yam, A., & Maoz, D. 2004, MNRAS, 347, 942
- Gehrels, N. 1986, ApJ, 303, 336
- Geldzahler, B. J., & Fomalont, E. B., 1984, AJ, 89, 1650
- Grggio, L. 2005, A&A, submitted
- Gregory, P.C., Scott, W.K., Douglas, K., & Condon, J.J. 1996, ApJS, 103,427
- Griffith, M. R., & Wright, A. E. 1993, AJ, 105, 1666
- Greggio, L., & Renzini, A. 1983, A&A, 118, 217
- Hamuy, M., Phillips, M.M., Maza, J., Suntzeff, N.B., Schommer, R.A., & Aviles, R. 1995,
AJ, 109,1
- Hamuy, M., Phillips, M. M., Suntzeff, N.B., Schommer, R. A.; Maza, J., & Aviles, R. 1996,
AJ, 112, 2391
- Hamuy, M., Trager, S. C., Pinto, P.A., Phillips, M. M., Schommer, R. A., Ivanov, V., &
Suntzeff, N.B. 2000, AJ, 120, 1479
- Hamuy, M., Phillips, M. M., Suntzeff, N. B. et al. 2003 Nature, 424, 651

- Heckman, T.M., Smith, E.P., Baum, S.A., van Breugel, W.J.M., Miley, G.K., Illingworth, G.D., Bothun, G.D., & Balick, B. 1986, *ApJ*, 311, 526
- Howell, D.A., 2001, *ApJ*, 554, L193
- Ivanov, V.D., Hamuy, M., & Pinto, P.A. 2000, *ApJ*, 542, 588
- Jarrett, T. H.; Chester, T.; Cutri, R.; Schneider, S. E.; & Huchra, J. P 2003, *AJ*, 125, 525
- Kennicutt, R.C., *ARA&A*, 36, 189
- Kochhar, R.K. 1989, *Astrophys. and Space Sci.*, 157, 305
- Kochhar, R.K., & Prabhu, T.P. 1984, *Astrophys. and Space Sci.*, 100, 369
- Ledlow, M.J., & Owen, F.N., 1996, *AJ*, 112, 9
- Li, W., Filippenko, A. V., Treffers, R. R.; Riess, A. G., Hu, J., & Qiu, Y. 2001, *ApJ*, 546, 734
- Livio, M., 2000, in *Type Ia Supernovae: Theory and Cosmology*, eds. J.C. Nyemeyer, & J.W. Truran (Cambridge: Cambridge Univ. Press), p. 33
- Livio, M., Riess, A., & Sparks, W., 2002, *ApJ*, 571, L99
- Livio, M., & Riess, A. 2003, *ApJ*, 594, L93
- Madau, P., Della Valle, M., & Panagia, N., 1998, *MNRAS*, 297, L17
- Mannucci, F., Della Valle, M., Panagia, N., Cappellaro, E., Cresci, G., Maiolino, R., Petrosian, A., & Turatto, M. 2005, *A&A*, in press (astro-ph/0411450)
- Maoz, D., & Gal-Yam, A. 2004, *MNRAS*, 347, 951
- Muriel, H. Nicotra, M. A., & Lambas, D.G. 1995, *AJ*, 110, 1032

- Panagia, N., 2000, in *Experimental Physics of Gravitational Waves*, eds. M. Barone, G. Calamai, M. Mazzoni, R. Stanga and F. Vetrano, (Singapore: World Scientific Publishing Co.), p. 107.
- Parodi, B.R., Saha, A., Sandage, A., & Tammann, G. A. 2000, *ApJ*, 540, 634
- Phillips, M. M., Jenkins, C. R., Dopita, M. A., Sadler, E. M., & Binette, L. 1986, *AJ*, 91, 1062
- Phillips, M.M. 1993, *ApJ*, 413, L105
- Phillips, M.M., Lira, P., Suntzeff, N.B., Schommer, R.A., Hamuy, M., & Maza, J. 1999, *AJ*, 118, 1766
- Prialnik, T., & Kovetz, A. 1995, *ApJ*, 445, 789
- Renzini, A., Ciotti, L., D’Ercole, A., & Pellegrini, S. 1993, *ApJ*, 419, 52
- Ruiz-Lapuente, P., Burkert, A., & Canal, R. 1995, *ApJ*, 447, L69
- Sadler, E.M., Jenkins, C.R., & Kotanyi C.G. 1989, *MNRAS*, 240, 591
- Sandage, A. 1986, *A&A*, 161, 89
- Srianand, R., & Gopal-Krishna 1998, *A&A*, 334, 39
- Strolger et al. 2004, *ApJ*, 613, 200
- Tinsley, B.M. 1974, *A&A*, 31, 463
- Tutukov, A.V., & Yungelson, L. 1994, *MNRAS*, 268, 871
- Tutukov, A.V., & Yungelson, L. 1996, *MNRAS*, 280, 1035
- Urry, C.M., & Padovani, P. 1995, *PASP*, 107, 803

van den Bergh, S., & Padzer J., ApJ, 390, 34

Whitmore, B.C., Zhang, Q. Leitherer, C., Fall, S. M., Schweizer, F., & Miller, B.W. 1999,
AJ, 118, 1551

Williams, R.E. 1994, ApJ, 426, 279

6. Captions

Fig. 1 Distribution of early-type galaxies in the Cappellaro et al. (1999) sample as a function of their absolute B-band magnitudes. The solid line is the distribution of the entire sample, the shaded histogram is the distribution of radio-loud galaxies, and the dashed histogram is the general luminosity function of elliptical galaxies, adapted from Muriel et al.(1995).

Fig. 2 Location of SN 1991bg and SN 1957B in NGC 4374. The Image is a collage of pictures of the galaxy and jets obtained from Digitalized Sky Survey (DDS) and NASA/IAPC Extragalactic Database (NED), respectively.


Fig. 3 Location of SN 1980N and SN 1981D in NGC 1316. Collage of images obtained from Digitalized Sky Survey (DDS) and NASA/IAPC Extragalactic Database (NED), respectively.

Fig. 4 Location of SN 1986G in NGC 5128. Collage of images obtained from Digitalized Sky Survey (DDS) and NASA/IAPC Extragalactic Database (NED), respectively.

Fig. 5 The rates of SNIa explosions (normalized to $10^{10} M_{\odot}$) as a function of the morphological Hubble type of the respective parent galaxies. Open symbols at $T \sim -2$ represent the SNIa rates of radio-loud (top) and radio-quiet (bottom) early-type galaxies. The filled symbol is the overall rate of SNIa events in early type galaxies, regardless of their radio properties (i.e. average of radio -loud and -quiet rates)

Fig. 6 The distribution of the light curve rates of decline for late (shade-dotted histogram) and early type galaxies (solid histogram) and the respective cumulative

distributions. The dashed box indicates the rates of decline for 8 objects belonging to the sample listed in Tab. 2.



This figure "f2.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0504087v1>

This figure "f3.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0504087v1>

This figure "f4.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0504087v1>

